



UDC 630; DOI 10.18551/rjoas.2022-10.06

## **EMPIRICAL STUDY OF FOREST REGENERATION POTENTIALITY OF NATURAL FOREST IN KADUNA NORTHERN GUINEA SAVANNAH: A CASE STUDY OF AFAKA FOREST RESERVE, NIGERIA**

**Sodimu A.I.\***

Savanna Forestry Research Station, Forestry Research Institute of Nigeria, Kaduna, Nigeria

**David B.**

Montane Forestry Research Station, Forestry Research Institute of Nigeria, Jos, Nigeria

**Adamu I.**

Shelterbelt Research Station, Forestry Research Institute of Nigeria, Kano, Nigeria

\*E-mail: [akintundesodimu@yahoo.com](mailto:akintundesodimu@yahoo.com)

### **ABSTRACT**

Forest Reserve Areas can sustain themselves of their valuable species through an effective regeneration system. At present, most of the preserved and permanent protected forest areas are been vandalized and valuable species are illegally removed. Based on the ecological and anthropogenic problems which lead to the extinction of valuable flora and fauna species, it became imperative to study species diversity and their potentiality for regeneration in natural scenario of Afaka Forest Reserve. Four (4) plots (A–D) of 100m x 100m were laid in each plot. Point centered quadrant method of sampling was used. Transect were selected on pure random basis where data were collected on species diversity and families, density of woody stem, diameter class distribution, regeneration potentials and relative frequency. 21 families with a total of 181 species were obtained on the species diversity at natural protected areas of Afaka Forest Reserve. The densities of woody plant varied between 625.01 - 961.51 per hectare in plots I, II, III and IV. While the highest number of trees in diameter-class distribution were apportioned to 10 – 19cm class interval in plots I, II, III and IV respectively. Diameter class 50cm and above had the lowest number of trees allocated to it. The regeneration potential of the diversified species was very poor, which has a great implication for regeneration and conservation of the various species encountered. Although *Isoberlinia doka* Craib & Stapf had the highest regeneration potential from each of the examined plots followed closely by *Acacia senegal*, *Parkia biglobosa* and *Terminalia avicennoides* Gull & Peer. Important woody plants of economic importance, like *Triplochiton scleroxylon* k. schum had zero regeneration potential. However, it is necessary to understand the phenology of the forest reserve. To study whether seeds or fruits produced are under adequate physiological conditions to germinate and grow into wildlings for regeneration purposes.

### **KEY WORDS**

Natural forest, forest regeneration, conservation, potentiality, vandalized.

Biological diversity conservation strategies that include a variety of individual management strategies for specific species are neither feasible nor effective (Attiwill, 1994; Borowski, 2015; Maggaji *et al.*, 2018). As an alternative, 'coarse-filter' management strategies focus on maintenance of broad patterns of forest age and composition under the assumption that this will provide the necessary habitat to support a wider variety of native species (Franklin, 1993, Larsen *et al.*, 1997). Native species are assumed to be adapted to the natural disturbance regime of their environment; thus, management that stimulates the natural disturbance regime has become a popular approach to maintaining species diversity as well as ecological integrity (Hunter, 1993, Bergeron *et al.*, 2002). With this approach, natural processes are to be protected, which include forest regeneration after disturbance



such as fire or logging (Sobamowo *et.al.*,2019). A central component of biodiversity conservation in managed forest landscape is successful regeneration of natural tree species mixes after logging (Bergeron and Harvey, 1997, Sodimu, *et. al.*,2020). This often proves to be a substantial challenge. Site preparation and planning of tree seedlings have been widely adopted as strategies to ensure rapid and successful postharvest regeneration. These require a substantial silvicultural and financial investment, however, and are not always successful in meeting legal regeneration standards let alone achieving a natural species mix. Although natural regeneration can be much less expansive and lead to more natural species mixes, its success depends on availability of seed sources, regeneration micro sites and micro environmental and biotic conditions favoring establishment and early survival of tree seedlings (Ngyuen – Xuan *et. al.*, 2000; Ogunkalu *et. al.*,2017).

There is necessity to conserve the forest resources because of their economic importance. Several direct as well as indirect benefits are available in the forest such as timber and non-timber products. Apart from timber products in the forest, plant foods such as leaves, seed nuts, fruits, tubers and roots stand as source of food income generation (Oduwaiye *et. al.*, 2002; Sodimu *et. al.*,2020). Also, species of economic importance such as medicinal and aesthetic values can be kept in perpetuity in a well preserved forest. The forests of Nigeria comprise of swamp forest, tropical rainforests and secondary forest regrowth. The lowland tropical rainforest was described by as a complicated mosaic of communities of different status and floristic composition (Sodimu, *et. al.*,2019). Structurally, the forests consist of three layers – the tree layer, the shrub layer and the undergrowth. Many of the emergent species in the forest grow slowly and require fair amount of overhead shade in their early life in order to survive and thrive (Dinenda *et.al.* 2015; Sodimu, *et. al.*, 2019) But it is now recognized that many of them grow very rapidly in full light conditions (Areola, 1991;).

The emergent species exist for a long period in a state of suppressed regrowth under shade of the canopy until a gap occurs which allows them to grow up. Forestry growth cycle consists of a mosaic gap-phase, build up phase and mature phase forests as explained by Van der maarel (1996). At present most of the preserved forest areas are being vandalized and valuable species illegally removed. The only mean by which Forest Reserved Areas can sustain themselves of these valuable species, is through an effective regeneration programmes. NEST (1991) indicated that 350,000 ha of forest and natural vegetation were lost in Nigeria. It led to problems such as desertification, soil erosion, declining soil fertility; flooding and extinction of important flora and fauna species (IUCN, 1980; Gabriel and Adedapo,2017). Forests are dynamic, seedlings germinate, grow, and compete with each other and with large trees. Some survive for hundreds of years. Changes will happen which will be predominant in the future forest depends not only on climate and soils but also on management decisions made today. Changes in forest composition will affect the quality and variety of forest resources available to future generations and wildlife (Sodimu *et. al.*2019). Sustainable management of forest ecosystem is set as a goal by most nations (Norton, 1996; Dinendra *et. al.*, 2015), and maintaining biological diversity is one important component of this (Hansen *et. al.*, 1991).

High demands for timber and increasing accessibility are leading to increased exploitation of these high elevation forests. They are characterized by slow growth rates of post-disturbance regeneration and established trees. Thus meeting objections for regeneration of natural species mixes and sustainability of fiber and other ecosystem components can prove challenging (Selmants and Knight, 2003; Sodimu *et. al.*, 2017). The only mean by which Forest Reserved Areas can sustain themselves of these valuable species, is through an effective regeneration programmes. Forest and natural vegetation lost in Nigeria every year due to anthropogenic activities are of great concerned. It led to problems such as desertification, soil erosion, declining soil fertility, flooding and extinction of important flora and fauna species (IUCN, 1980). Hence, necessitate the need to study the natural regeneration potentials of forest reserved areas so that species of economic significance, including non-timber forest products (NTFPs) and aesthetic values, can be



maintained in perpetuity. The objective of the study is to assess the species diversity and potentiality for forest regeneration in natural forest of Afaka Forest Reserve.

## METHODS OF RESEARCH

The study was carried out in natural forest of Afaka Forest Reserve. Afaka is in Igabi Local Government Area. Igabi Local Government Area is located in the Northern Guinea Savanna region of Nigeria on latitude  $10^{\circ}E$   $37^{\circ}N$  /  $10^{\circ}$   $41^{\circ}N$  and longitude  $7^{\circ}47^{\circ}E$  /  $7^{\circ}51^{\circ}E$  (Otegbeye *et. al.*, 2001) Igabi local Government Area shares boundary with Kaduna South, Kaduna North, Giwa Local Government and Zaria Local Government areas of Kaduna State. It has an Annual rainfall of about 1000mm -1500mm per annum. Major crops grown in the area include maize, cassava, millet, sorghum, guinea-corn and water-melon. The area consists of different tribes and ethnic groups such as Yoruba's, Hausa, Igbo's, Fulani's, Gbagyi's and Hausa's. The area has an estimated population of about 570,00 people and covers an area of about 4556.95 square kilometers. (NPC,2006)

Four (4) sampling plots of 100m x 100m were laid in Old Afaka Forest Reserve. Technique used was adopted from Cuttam and Curts, 1956; Oduwaiye *et. al.*,2002; Sodimu *et.al*, 2017. Point centred quadrant method of sampling was used in each of the selected plots. Transects were selected using systematic sampling method. The first one was selected on pure random basis. Along the direction of each transect, sampling points were taken at intervals of 5m which helps in getting sufficient sampling points and possible overlapping and duplication of units of measurement were avoided (Sodimu, *et. al.*,2020) At each sampling point, the following parameters were assessed. Distance to four nearest woody plants, one in such quadrant.

Height of the trees measured through the use of Haga altimeter.

Diameter of each tree measured through the use of diameter tape.

Densities of the woody stem in all the study plots were determined. The distance measured between sample points of each woody plant was used to obtain the density stem. The estimation was based on the formula:

$$D = \frac{10^4 \text{ stem ha}^{-1}}{d^2}$$

Where:  $d^2$  = is the average distance in  $m^2$  (Cuttam and Curtis 1956; Sodimu, *et. al.*, 2020).

Diameter class distribution was determined through the measurement of diameter of the woody stand in all the plots at DBH (1.37m height).

Regeneration potentials:

$$\text{Regeneration potential} = \frac{\text{Number of wilding of individual species}}{\text{Density of the woody stem}}$$

Relative frequency:

$$R.F = \frac{\text{Number of individual of the species}}{\text{Number of individual of all species}} \times 100$$

Species abundance: indicating the number of species contained in the number of individual plants.

## RESULTS AND DISCUSSION

Members of the families in acceding order include *Fabaceae*>*Combretaceae*, *Sterculiaceae*>*Verbanaceae* dominated the permanent protected area followed by members of *Annonaceae* >*Rhamnaceae*> *Anacardiaceae* >*Meliaceae* and to a lesser extent, some



other families include *Sapotaceae*>*Euphorbiaceae*> *Ochnaceae* and so on. The species that are most common in acceding order are *Isobanilia doka*>*Parkia biglobosa*>*Prosopis africana*>*Vitex doniana*> *Acacia senegal* > *Lawsonia inermis* (Table 1). These results are in agreement with the work of Sodimu *et. al.* (2020) that the first four (4) families dominated Old Afaka and Buruku forest Reserve with high percentages spread.

Table 1 – Species Diversity and Families in Natural Forest of Afaka Forest Reserve

S/No	Families	Number of Species	Percentage (%) Spread
1	Fabaceae	28	15.4
2	Ebenaceae	6	3.32
3	Annonaceae	8	4.42
4	Anacardiaceae	7	3.87
5	Combretaceae	15	8.28
6	Longaniaceae	5	2.76
7	Meliaceae	7	3.87
8	Rhamnaceae	7	3.70
9	Rubiaceae	6	3.87
10	Sapotaceae	8	4.42
11	Sterculiaceae	10	5.53
12	Verbanaceae	7	3.87
13	Papilioceae	8	4.42
14	Casalpiniaceae	6	3.32
15	Myristicaceae	7	3.87
16	Ochnaceae	8	4.42
17	Irvingiaceae	10	5.53
18	Chysobaniaceae	6	3.32
19	Euphorbiaceae	8	4.42
20	Lythraceae	8	4.42
21	Myrtaceae	6	3.32
	TOTAL	181	100

The density of the woody stem as indicated in table 2, revealed that plot C in permanent protected area had 961.51stem/ha which was the highest when compare to plot A, plot B and plot D with 857.64stem/ha, 760.72stem/ha and 625.01stem/ha respectively.

Table 2 – Density of Woody Plants at Various Plots in Natural Forest of Afaka Forest Reserve

S/N	Site	Stem (ha <sup>-1</sup> )
1	Plot III	961.50
2	plot I	857.64
3	Plot II	760.72
4	Plot IV	625.01

Diameter class distribution of woody stems assessed in plot A, B, C and D; however, the number of trees assessed for diameter was 80, 125, 85 and 80 respectively. The highest number of trees was apportioned to diameter class 10 – 19cm for each of the plots A, B, C and D, that was 40 trees, 60 trees, 75 trees and 40 trees respectively, followed by diameter class 20 – 29cm (fig. 1). Diameter class 50cm and above had the lowest number of trees allocated to it. Perhaps, this might be due to over exploitation and activities of illegal fellers in the reserve hunting for trees with bigger girth for conversion into sizeable wood. The above observation was in agreement with the work of Sodimu, *et. al.*, (2020) who also noted that diameter class of trees between 10cm-19cm are commonly found in most forest reserves around Kaduna Northern Guinea Savanna eco-zones.

The regeneration potential of the diversified species as indicated in table 3 is very poor. It therefore, has a great implication for regeneration and conservation of the various species encountered. *Isobanilia doka* had the highest regeneration potential from each of the examined plots in Natural Forest of Afaka Forest Reserve; followed by *Acacia senegal*, *Parkia biglobosa* and *Terminalia avicennoides*. However, most of the important woody plants of economic importance like *Triplochiton scleroxylon*, had zero regeneration potential. Regeneration potential of the most species is zero, owing to complete absent of mother trees which would have produced the wildlings within the assessed area. The results are in



consonance with the works and observations of Oduwaiye *et. al.*, (2002); Sodimu *et. al.*, (2017); Sodimu *et. al.*, (2020).




Figure 1 – Diameter Class distribution of woody stem in plot I, II, III and IV of Natural Forest of Afaka Forest Reserve

Table 3 – Regeneration Potential of Species Assessed in Natural Forest of Afaka Forest Reserve

S/N	Species	Plot I	Plot II	Plot III	Plot IV
1	<i>Prosopis africana</i>	0.00	0.00	0.00	0.03
2	<i>Entada Africana</i> Gull & Perr.	0.01	0.00	0.00	0.00
3	<i>Vitex doniana</i> Linn.	0.00	0.00	0.00	0.01
4	<i>Terminalia avicennoides</i> Guill & Perr.	0.05	0.02	0.01	0.02
5	<i>Macaranga hurifolia</i>	0.00	0.00	0.00	0.00
6	<i>Triplochiton scleroxylone</i> K.Schum	0.00	0.00	0.00	0.00
7	<i>Isobertinia doka</i> Craib & Stapf.	0.44	0.75	0.76	0.68
8	<i>Annona senegalensis</i> Pers.	0.00	0.05	0.00	0.00
9	<i>Carissa adolia</i> Linn.	0.01	0.00	0.00	0.00
10	<i>Pilostigma thornningii</i> (schum)	0.00	0.00	0.03	0.00
11	<i>Acacia senegal</i> (Houtt)	0.03	0.02	0.04	0.01
12	<i>Lophira lanceolata</i> Banks ex Gaertn.	0.01	0.00	0.00	0.00
13	<i>Combretum ghasalense</i> Engl. & Diels	0.01	0.00	0.01	0.00
14	<i>Parkia biglobosa</i> (Jacq) Benth	0.02	0.04	0.01	0.02
15	<i>Parinari curatellifolia</i> Planch.	0.00	0.02	0.00	0.00

Table 4 – Relative Frequency of Species in Natural Forest of Afaka Forest Reserve

S/N	Species	Plot I	Plot II	Plot III	Plot IV
1	<i>Terminalia avicennoides</i> Guill & Perr.	5.850	8.492	6.462	8.739
2	<i>Khaya senegalensis</i> Ders.	0.278	0.450	0.303	0.000
3	<i>Vitellaria paradoxa</i> C.F. Gaertn	0.543	1.046	1.256	1.817
4	<i>Pilostigma thornningii</i> (schum)	4.403	2.063	5.229	2.008
5	<i>Annona senegalensis</i> Pers.	8.951	3.487	7.757	0.000
6	<i>Acacia Senegal</i> (Houtt)	67.208	56.010	46.644	45.372
7	<i>Lannea barteri</i> (Oliv)	1.733	0.699	0.000	1.205
8	<i>Gardenia imperialis</i> (K.Schum)	1.338	5.846	0.000	1.012
9	<i>Carissa adolia</i> Linn.	4.703	1.848	5.687	2.460
10	<i>Sterculia setigera</i> Del.	1.538	7.946	0.000	6.422
11	<i>Isobertinia doka</i> Craib & Stapf.	8.188	20.541	7.653	14.879
12	<i>Entada Africana</i> Gull & Perr.	0.349	0.320	0.000	0.904
13	<i>Diospyros mespiliformis</i> Hochst.	0.368	0.000	0.000	0.304
14	<i>Prosopis Africana</i> (Gull & Perr) Taub	0.412	0.000	0.000	0.000
15	<i>Combretum ghasalens</i> Engl. & Diels	0.288	1.139	0.000	0.000
16	<i>Anogeisus liocarpus</i> (DC) Gull & Perr	6.850	0.000	0.000	0.501
17	<i>Parkia biglobosa</i> (Jacq) Benth	5.465	0.000	0.00	0.000
18	<i>Triplochiton scleroxylon</i> K.Schum.	0.890	0.000	1.711	0.000
19	<i>Vitex doniana</i> Linn	0.890	0.000	1.811	1.306
20	<i>Strychnos spinosa</i> Lam.	0.418	0.000	2.005	0.000
21	<i>Tectona grandis</i> L.f.	2.256	0.000	0.000	1.605
22	<i>Macaranga hurifolia</i> Beille.	0.000	9.692	1.247	6.745
23	<i>Parinari curitellifolia</i> Planch.	0.000	0.460	0.000	0.604
24	<i>Lophira lanceolata</i> Banks ex Gaertn	0.000	0.560	2.467	0.000
25	<i>Lawsonia inermis</i> L.	0.000	0.256	7.242	0.000
26	<i>Psidium guajava</i> L.	0.000	0.350	0.303	1.705
27	<i>Irvingia gabonensis</i> (Bail)	0.000	1.089	1.543	1.309





In table 4 plots A, B, C and D, 21, 19, 16 and 18 species were observed. In plot A, *Acacia senegal* showed the greatest occurrence with relative frequency of 67.2%, the species had some spread in plot B, plot C and plot D with relative frequency of 56%, 46.6% and 45.4%. This was followed by *Isoberlinia doka* in plots B and plot D with relative frequency of 20.5% and 15.9% respectively. However, in plot C *Acacia senegal* was closely followed by *Annona senegalensis* with relative frequency of 7.56% respectively. This is in agreement with the work of Sodimu, *et. al.*, 2020 who observed that *I. doka* and *A. senegal* have better spread in Northern Guinea Savannah eco-zone.

### CONCLUSION

None of the tree species in protected area of natural forest of Afaka Forest Reserve was observed flowering during the period of assessment. However, it could not be ascertained whether most of the diverse species had undergone generative phase or were still undergoing vegetative phase. From the analysis, it would be observed that most of the woody species were not regenerating themselves. There were cases when mother trees of the wildlings observed were absent from the plots or the wildlings of woody species were absent. This situation poses a great threat to conservation of the diverse species in the Forest Reserve. Based on the findings, it is recommended that it is of paramount importance to understand the phenology of the forest reserve and, to study whether seed or fruits produced are under adequate physiological conditions to germinate and grow into wildlings for regeneration purpose. Otherwise, it will be very necessary to find means of regenerating the reserve either artificially or naturally for sustainability of the forest reserve.

### REFERENCES

1. Areola, O. E (1991). Ecology of Natural Resources in Nigeria. CAB.; 278.
2. Attiwill, P.M. (1991). The Disturbance of Forest Ecosystems: The Ecological Basis for Conservative Management. *For. Ecol. Manag.* 63:247–300.
3. Bergeron Y, A. Leduc, B. Harvey, and S. Gauthier. (2002) A Guide for Sustainable Management of the Canadian Boreal Forest. *Silva. Fenn.* 36:81–95.
4. Bergeron, Y., and B. Harvey. (1997) Basing Silviculture on Natural Ecosystem Dynamics: An Approach Applied to the Southern Boreal Mixed Wood Forest of Quebec. *For. Ecol. Manag.* 92: 235–242.
5. Borowski, T (2015). Modifications of Natural Rubber (*Hevea brasiliensis*): Production, Application and Comparison. *International journal of World Scientific News*, 20-5.
6. Cuttam, G. and Curtis J.T. (1956). Use of Distance Measure in Phytosociological Sampling. *Ecology*, 37, 451- 460.
7. Dinendra Raychaudhuri, Sumana Saha, Souvik Sen, Dhruva Chandra Dhali (2015). Spiders (Araneae: Arachnida) of Reserve Forests of Dooars: Gorumara National Park, Chapramari Wildlife Sanctuary and Mahananda Wildlife Sanctuary. *International Journal of World Scientific News.*;20:1-339.
8. Franklin, J.F (1993) Preserving biodiversity Species, Ecosystems, or Landscapes? *Ecol. Appl.* 34: 202–205.
9. Gabriel, A.A and Adedapo, A.A(2017). Exotic Versus Indigenous and Implication for Environmental Forestry Management in the Niger Delta, Nigeria. *International Journal of World Scientific News.*;74:36.
10. Hansen, A.J., T.A. Spies, and F.J. Swanson. (1991) Conserving Biodiversity in managed Forests Lessons from Natural Forests. *Bioscience.* 41:382–392.
11. Hunter, M. L. (1993) Natural fire regimes as spatial models for managing boreal forests. *Biol. Conserv.* 65:115–120.
12. IUCN World Conservation Strategy (1980). Living Resources conservation for Sustainable Development -IUCN-UNEP-WWF 1980.



13. Larsen, D.L., R. Shiftly, F.R. Thomson, B.L. Bokkshire, D.C. Dey, E.W. Kurzejeski, and K. England. (1997). Ten Guidelines for Ecosystem Researchers: Lessons from Missouri. *J. For.* 95: 4-8.
14. Magaii, Y., Ajibade, G.A., Yilwa, V.M., Appah, J., Haroun, A.A., Alhaji, I., Umadi, M.M. and Sodimu, A.I. (2019). Concentration of Heavy Metals in the Soil and Translocation with Phytoremediation Potential by Plant species in Military Shooting Range. *International Journal of World Scientific News*. Vol. 92(3):260-271. EISSN: 2392 – 2192.
15. National Population Commission (2006). National Population Commission (N.P.C.) News Bulletin, Kaduna State. 13-15.
16. Nguyen-Xuan, T., Y. Bergeron, D. Simard, J.W. Fyles, and D. Pare. (2000) The importance of Forest Floor Disturbance in the Early Regeneration Patterns of the Boreal Forest of Western and Central Quebec: A Wildfire Versus Logging Comparison. *Can. J. For. Res.* 30:1353–1364.
17. Nigerian Environmental Study Team Nigeria's Threatened Environment (1991) A National Research Profile. Nigerian Environmental Study/Action Team (NEST), Ibadan, Nigeria. 287 – 290.
18. Norton, T.W (1996) Conservation of biological diversity in temperate and boreal forest ecosystems. *For. Ecol. Manag.* 85:1–7.
19. Oduwaiye, E.A. Oyeleye, B and Oguntala, A.B (2002). Species diversity and potentiality for forest regeneration in Okomu permanent sample plot. *Proceeding of the 28th Annual conference of the Forestry Association of Nigeria*. 264-269.
20. Ogunkalu, O.A., Sodimu, A.I., Suleiman, R.A. and Adedire, O.O. (2017). Survey of benefits and Constraints of Urban Trees in Kaduna Metropolis. *International Journal of World News of Natural Sciences*, Vol. II Pp 19-27. EISSN: 2543-5426. [www.woldnewsnaturalsciences.com](http://www.woldnewsnaturalsciences.com).
21. Selmants, P.C., and Knight, D.H. (2003). Understory Plant Species Composition 30–50 years After Clear Cutting in Southeastern Wyoming Coniferous Forests. *Forest Ecology Management* 185:275–289.
22. Sobamowo, G, Kamiyo, O.M and Adeoye, O.A. (2019) Temporal Variation in Species Composition Diversity and Regeneration Status Along Altitudinal Gradient Slope. *International Journal of World Scientific News*. 138:2:167-191.
23. Sodimu, A.I., Onwumere, G.B. and Yilwa, V.M. (2019). Assessment of Soil Based Heavy Metals from Anthropogenic Activities in Kaduna Guinea Savanna of Nigeria. *International Journal of World Scientific News*. Vol. 125: 83-93 EISSN: 2392 – 2192.
24. Sodimu, A.I., Yilwa, V.M. and Onwumere. G.B. (2017). The Impact of Gas Flaring from Kaduna Refining and Petrochemical Company (KRPC) on Plant Diversity in Kaduna Northern Guinea Savanna Eco-Region of Nigeria. *International Journal of World Scientific News*. Vol.69:168 – 178. EISSN: 2392 – 2192.
25. Sodimu, A.I., Ogunkalu, O.A., Komolafe, A.S., and Sadiq, Z.K. (2017). Economic Analysis of Wonderful Kola (*Hydrocotyle asiata*) Marketing in Kaduna North Local Government Area of Kaduna State, Nigeria. *International Journal of World News of Natural Sciences*. Vol. 9, PP 1 – 6. EISSN: 2543 – 5426.
26. Sodimu, A.I., Onwumere, G.B. and Yilwa, V.M. (2019). Soil Based Heavy Metals Originating from Anthropogenic Activities on Floristic Composition of Some Selected Sites in Kaduna Northern Guinea Savanna of Nigeria. *Asian Journal of Environment and Ecology*. Vol. 10 (4); 1-10. ISSN: 2456 – 690X. DOI: 10.9734/AJEE/2019/V10: 430122.
27. Sodimu, A.I, Sulaiman, R.A, Maikano, S and Lapkat, G.L (2020). Assessment of Species Diversity and Potentiality in Buruku Forest Reserve, Kaduna State, Nigeria. *Asian Journal of Research in Agriculture and Forestry*. 5(1): 39-46. ISSN: 2581-7418. DOI: 9734/AJRAF/2020/v5i130077.
28. Van Der Maarel, E (1996) Pattern and Process in Plant Community: Fifty Year After A. S. Watt. *Journal of Vegetation Science* 7: 19- 28.